

8-2012

# Creating a model antigen system to test the mechanism of GCC-specific tolerance

Patrick Ihejirika

Lincoln University, Patrick.ihejirika@lincoln.edu

## Let us know how access to this document benefits you

Follow this and additional works at: <http://jdc.jefferson.edu/summercancerimmunotherapy>Part of the [Pharmacy and Pharmaceutical Sciences Commons](#)


### Recommended Citation

Ihejirika, Patrick, "Creating a model antigen system to test the mechanism of GCC-specific tolerance" (2012). *Summer Training Program in Cancer Immunotherapy*. Presentation 4.  
<http://jdc.jefferson.edu/summercancerimmunotherapy/4>

This Article is brought to you for free and open access by the Jefferson Digital Commons. The Jefferson Digital Commons is a service of Thomas Jefferson University's [Center for Teaching and Learning \(CTL\)](#). The Commons is a showcase for Jefferson books and journals, peer-reviewed scholarly publications, unique historical collections from the University archives, and teaching tools. The Jefferson Digital Commons allows researchers and interested readers anywhere in the world to learn about and keep up to date with Jefferson scholarship. This article has been accepted for inclusion in Summer Training Program in Cancer Immunotherapy by an authorized administrator of the Jefferson Digital Commons. For more information, please contact: [JeffersonDigitalCommons@jefferson.edu](mailto:JeffersonDigitalCommons@jefferson.edu).

# CREATING A MODEL ANTIGEN SYSTEM TO TEST THE MECHANISM OF GCC-SPECIFIC TOLERANCE

○ Patrick Ihejirika



# Cancer Cases and Deaths

## Leading New Cancer Cases and Deaths – 2012 Estimates

Estimated New Cases*		Estimated Deaths	
Male	Female	Male	Female
Prostate 241,740 (29%)	Breast 226,870 (29%)	Lung & bronchus 87,750 (29%)	Lung & bronchus 72,590 (26%)
Lung & bronchus 116,470 (14%)	Lung & bronchus 109,690 (14%)	Prostate 28,170 (9%)	Breast 39,510 (14%)
Colon & rectum 73,420 (9%)	Colon & rectum 70,040 (9%)	Colon & rectum 26,470 (9%)	Colon & rectum 25,220 (9%)
Urinary bladder 55,600 (7%)	Uterine corpus 47,130 (6%)	Pancreas 18,850 (6%)	Pancreas 18,540 (7%)
Melanoma of the skin 44,250 (5%)	Thyroid 43,210 (5%)	Liver & intrahepatic bile duct 13,980 (5%)	Ovary 15,500 (6%)
Kidney & renal pelvis 40,250 (5%)	Melanoma of the skin 32,000 (4%)	Leukemia 13,500 (4%)	Leukemia 10,040 (4%)
Non-Hodgkin lymphoma 38,160 (4%)	Non-Hodgkin lymphoma 31,970 (4%)	Esophagus 12,040 (4%)	Non-Hodgkin lymphoma 8,620 (3%)
Oral cavity & pharynx 28,540 (3%)	Kidney & renal pelvis 24,520 (3%)	Urinary bladder 10,510 (3%)	Uterine corpus 8,010 (3%)
Leukemia 26,830 (3%)	Ovary 22,280 (3%)	Non-Hodgkin lymphoma 10,320 (3%)	Liver & intrahepatic bile duct 6,570 (2%)
Pancreas 22,090 (3%)	Pancreas 21,830 (3%)	Kidney & renal pelvis 8,650 (3%)	Brain & other nervous system 5,980 (2%)
All sites 848,170 (100%)	All sites 790,740 (100%)	All sites 301,820 (100%)	All sites 275,370 (100%)

\*Excludes basal and squamous cell skin cancers and in situ carcinoma except urinary bladder.

# Colorectal Cancer

- The main treatment options include:  
Surgery, chemo-, radiation, biological therapy.  
**All treatments present a risk of side effects**
- Current standard of care: FOLFOX, FOLFIRI, etc
  - ✓ *Developed in 1980*
  - ✓ *Minimal response rate*
  - ✓ *Prognosis: 5 year survival rate of 63%*

# Colorectal Cancer Immunotherapy

***Meta-analysis of clinical trials reveal a very weak clinical response rate (<1%) for active specific immunotherapy procedures currently available for colorectal cancer***

Nagorsen D, et al. Clin Cancer Res 2006;12:3064-9.


***Failure is due to the lack of molecularly defined tumor-associated antigens that can be reliably considered:***

- Tumor-specific***
- Sufficiently immunogenic***
- Shared among different patients***

Adapted from: Dalerba P, et al. Crit Rev Oncol Hemat 2003;46(1):33-57.

Colorectal Cancer TAA		
<u>Mutant Self Proteins</u>		
K-ras	p53	
<u>Oncofetal / Cancer Testis Antigens</u>		
βhCG	Gastrin	5T4
<u>Overexpressed Self Antigens</u>		
p53	MUC1	SART
Sialyl-Tn	Her2/neu	ART
Survivin	CD55	Ep-CAM
Carcinoembryonic Antigen (CEA)		
<u>Tissue-Specific Differentiation Antigens</u>		
	?	



# Guanylyl Cyclases



Guanylyl Cyclases		
	Isoforms	Ligands
Natriuretic Peptide Receptors	GCA	ANP, BNP
	GCB	CNP
Intestinal Peptide Receptor	GCC	ST
		Guanylin Uroguanylin
Orphan Receptors	GCD	
	GCE	
	GCF	
	GCG	
Soluble	sGC	NO




# Mucosa-Restricted GCC Expression



Swensen et al. 1996. Biochemical and Biophysical Research Communications, 225: 1009-1014

S Schulz et al, unpublished data

E Lin et al, unpublished data








# Anti-Tumor Immunity

Control AV

GCC-AV




Control AV

GCC-AV







# GCC-Specific Immune Responses







# Predicted CD4 Tolerance Mechanism








# Model System



# Tac





6μm slice





# Tac FACS

Extracellular





Intracellular





# Epitopes and surface expression




# Creating GCC-based model antigen



## Steps:

- cloning different epitopes and signal sequences into GCC:
  - ✓ Bip and Prp
- clone the constructs into mammalian expression plasmid (retrovirus)
- create cell lines expressing the constructs by infecting them with retrovirus containing our constructs
- test the new cell lines for 1) Chimeric antigen expression, 2) Chimeric antigen subcellular location, 3) epitope presentation
- work with Dr Eisenlohr's lab to see how different signal sequences and model epitopes affect our new model antigen.



# Overhang PCR



# PCR Epitopes and GCC




# Overhang PCR





# Screening pENTR clones



# Cutting inserts out

- Sequenced pENTR clones
- Cut inserts out with NotI and SalI
- Cut pMSCV-Puro with NotI and SalI
- Ligated inserts with pMSCV-Puro
- Transformed bacteria
- Screened colonies

# Results



# Summary/Future Goal

- Create stable cell lines expressing the constructs by target cell lines with retrovirus containing our constructs.
- Test the new cell lines for 1) chimeric antigen expression, 2) define the subcellular location of our antigen, 3) quantify epitope presentation
- Work with Dr. Eisenlohr's lab to see how different signal sequences and model epitopes affect our new model antigen.

# Thank You!

- Thomas Jefferson University:
- Elizabeth Rappaport MD
- Scott Waldman MD/PhD
- Adam Snook PhD
- Michael Magee
- Bo Xiang
- Laurence Eisenlohr VMD, PhD
- St. Joseph University
- Audrey Fritzinger